arXiv:math/0410529v2 [math.CO] 8 Jun 2005

Final version for J. Number Theory, `arXiv:math.CO/0410529`.

ON VARIOUS RESTRICTED SUMSETS

Zhi-Wei Sun[1,*] and Yeong-Nan Yeh[2]

[1]Department of Mathematics and Institute of Mathematical Science
Nanjing University, Nanjing 210093, P. R. China
zwsun@nju.edu.cn
`http://pweb.nju.edu.cn/zwsun`

[2]Institute of Mathematics, Academia Sinica, Taipei, Taiwan
mayeh@ccvax.sinica.edu.tw

Abstract. For finite subsets $A_1,\ldots,A_n$ of a field, their sumset is given by $\{a_1+\cdots+a_n:\ a_1\in A_1,\ldots,a_n\in A_n\}$. In this paper we study various restricted sumsets of $A_1,\ldots,A_n$ with restrictions of the following forms:

$$a_i-a_j\not\in S_{ij},\ \text{or}\ \alpha_i a_i\neq\alpha_j a_j,\ \text{or}\ a_i+b_i\not\equiv a_j+b_j\ (\mathrm{mod}\ m_{ij}).$$

Furthermore, we gain an insight into relations among recent results on this area obtained in quite different ways.

1. Introduction

The additive order of the identity of a field F is either infinite or a prime, we call it the *characteristic* of F.

Let F be a field of characteristic p, and let $A_1,\ldots,A_n$ be finite subsets of F with $0<k_1=|A_1|\leqslant\cdots\leqslant k_n=|A_n|$. Concerning various restricted sumsets of $A_1,\ldots,A_n$, the following results are known:

(i) (The Cauchy-Davenport theorem (see, e.g. [N]))

$$|\{a_1+\cdots+a_n:\ a_1\in A_1,\ldots,a_n\in A_n\}|\geqslant\min\{p,k_1+\cdots+k_n-n+1\}.$$

(ii) (Dias da Silva and Hamidoune [DH]) If $A_1=\cdots=A_n=A$, then

$$|\{a_1+\cdots+a_n:\ a_i\in A,\ a_1,\ldots,a_n\ \text{are distinct}\}|\geqslant\min\{p,n|A|-n^2+1\}.$$

2000 *Mathematics Subject Classification.* Primary 11B75; Secondary 05A05, 11C08.

[*]This author is responsible for communications, and supported by the National Science Fund for Distinguished Young Scholars (No. 10425103) and the Key Program of NSF (No. 10331020) in China.

(iii) (Alon, Nathanson and Ruzsa [ANR2]) If $k_1 < \cdots < k_n$, then

$$|\{a_1+\cdots+a_n : a_i \in A_i,\ a_i \neq a_j \text{ if } i \neq j\}| \geqslant \min\left\{p, \sum_{i=1}^n k_i - \frac{n(n+1)}{2} + 1\right\}.$$

(iv) (Hou and Sun [HS]) Let S_{ij} $(1 \leqslant i,j \leqslant n,\ i \neq j)$ be finite subsets of F with cardinality m. If $k_1 = \cdots = k_n = k$ and $p > \max\{ln, mn\}$ where $l = k-1-m(n-1)$, then

$$|\{a_1+\cdots+a_n : a_i \in A_i,\ a_i - a_j \notin S_{ij} \text{ if } i \neq j\}| \geqslant ln+1.$$

(v) (Liu and Sun [LS]) Let $P_1(x), \ldots, P_n(x) \in F[x]$ be monic and of degree $m > 0$. If $k_n > m(n-1)$, $k_{i+1} - k_i \in \{0,1\}$ for all $i = 1, \ldots, n-1$, and $p > K = (k_n-1)n - (m+1)\binom{n}{2}$, then we have

$$|\{a_1+\cdots+a_n : a_i \in A_i,\ P_i(a_i) \neq P_j(a_j) \text{ if } i \neq j\}| \geqslant K+1.$$

(vi) (Sun [Su]) Let $P_1(x), \ldots, P_n(x) \in F[x]$ have degree $m > 0$ with the permanent of the matrix $(b_j^{i-1})_{1\leqslant i,j\leqslant n}$ nonzero, where b_j is the leading coefficient of $P_j(x)$. If $k_1 = \cdots = k_n = k > m(n-1)$ and $K = (k-1)n - (m+1)\binom{n}{2} < p$, then

$$|\{a_1+\cdots+a_n : a_i \in A_i,\ a_i \neq a_j\ \&\ P_i(a_i) \neq P_j(a_j) \text{ if } i \neq j\}| \geqslant K+1.$$

While result (ii) was deduced by a deep tool from the representation theory of symmetric groups, results (iii)–(vi) were obtained from the following basic principle arising from Alon and Tarsi [AT].

Combinatorial Nullstellensatz ([A1, A3]). *Let $A_1, \ldots, A_n$ be finite subsets of a field F with $|A_i| > k_i$ for $i = 1, \ldots, n$ where $k_1, \ldots, k_n$ are nonnegative integers. If the coefficient of the monomial $x_1^{k_1} \cdots x_n^{k_n}$ in $f(x_1, \ldots, x_n) \in F[x_1, \ldots, x_n]$ is nonzero and $k_1+\cdots+k_n$ is the total degree of f, then there are $a_1 \in A_1, \ldots, a_n \in A_n$ such that $f(a_1, \ldots, a_n) \neq 0$.*

Lower bounds for various restricted sumsets are usually yielded with help of the following lemma (or Proposition 2.1 of [HS]) implied by the Combinatorial Nullstellensatz.

Lemma 1.1 (Alon et al. [ANR1, ANR2]). *Let $A_1, \ldots, A_n$ be finite nonempty subsets of a field F with $k_i = |A_i|$ for $i = 1, \ldots, n$. Let $P(x_1, \ldots, x_n) \in F[x_1, \ldots, x_n] \setminus \{0\}$ and $\deg P \leqslant \sum_{i=1}^n (k_i - 1)$. If the coefficient of the monomial $x_1^{k_1-1} \cdots x_n^{k_n-1}$ in the polynomial*

$$P(x_1, \ldots, x_n)(x_1+\cdots+x_n)^{\sum_{i=1}^n (k_i-1) - \deg P}$$

does not vanish, then we have

$$|\{a_1+\cdots+a_n : a_i\in A_i,\ P(a_1,\dots,a_n)\neq 0\}| \geqslant \sum_{i=1}^n (k_i-1)-\deg P+1.$$

In the next section, we will develop a general technique to compute certain coefficients of some polynomials. Using Lemma 1.1 and our work in Section 2, we will prove the following main theorems in Section 3.

Theorem 1.1. *Let F be a field of characteristic p, and let $A_1,\dots,A_n$ be finite nonempty subsets of F with $|A_n|=k$ and $|A_{i+1}|=|A_i|+1$ for $i=1,\dots,n-1$. Let m be a positive integer, and let $S_{ij}\subseteq F$ and $|S_{ij}|<2m$ for all $1\leqslant i<j\leqslant n$. If $p>\max\{mn,(k-1)n-mn(n-1)\}$, then we have*

$$|\{a_1+\cdots+a_n : a_i\in A_i,\ \text{and}\ a_i-a_j\notin S_{ij}\ \text{if}\ i<j\}| \geqslant (k-1-m(n-1))n+1.$$

Remark 1.1. Theorem 1.1 can be viewed as a partial generalization of result (iii).

Theorem 1.2. *Let k and m be positive integers. Let $A_1,\dots,A_n$ be subsets of the complex field $\mathbb{C}$ with cardinality k, and let S_{ij} $(1\leqslant i<j\leqslant n)$ be subsets of $\mathbb{C}$ with at most $2m-1$ elements. If $\zeta_1,\dots,\zeta_n$ are distinct qth roots of unity where q is a positive odd integer, then*

$$\left|\left\{\sum_{i=1}^n a_i : a_i\in A_i,\ a_i\zeta_i\neq a_j\zeta_j\ \text{and}\ a_i-a_j\notin S_{ij}\ \text{if}\ i<j\right\}\right| \geqslant (k-1-m(n-1))n+1.$$

Remark 1.2. A conjecture of Snevily [S] states that for any cyclic group with odd order if A and $B=\{b_1,\dots,b_n\}$ are its subsets with cardinality n then there is a numbering $\{a_i\}_{i=1}^n$ of the elements of A such that $a_1b_1,\dots,a_nb_n$ are pairwise distinct. Using the Combinatorial Nullstellensatz Alon [A1] confirmed this for the cyclic group $\mathbb{Z}/p\mathbb{Z}$ where p is an odd prime. Since we can identify a cyclic group of order q with the multiplicative group of all the qth roots of unity, Snevily's conjecture follows from Theorem 1.2 in the case $k=n$, $m=1$, $A_1=\cdots=A_n=A$ and $S_{ij}=\{0\}$ $(1\leqslant i<j\leqslant k)$, which was first obtained by Dasgupta, Károlyi, Serra and Szegedy [DKSS] in 2001. Another extension of Snevily's conjecture appeared in [Su].

Theorem 1.3. *Let $\alpha_1, \dots, \alpha_n$ be positive reals, and let $b_1, \dots, b_n$ be integers. Let $A_1, \dots, A_n$ be finite subsets of $\mathbb{Z}$ with cardinality $k > 0$. For $1 \leqslant i < j \leqslant n$ let m_{ij} be an integer greater than $2\max\{|x_i - x_j| : x_i \in A_i,\ x_j \in A_j\}|$. Then the restricted sumset*

$$\left\{ \sum_{i=1}^{n} a_i : a_i \in A_i,\ a_i\alpha_i \neq a_j\alpha_j \text{ and } a_i + b_i \not\equiv a_j + b_j \pmod{m_{ij}} \text{ if } i < j \right\}$$

has more than $(k-n)n$ elements.

Corollary 1.1 (Kézdy and Snevily [KS]). *Let m and n be positive integers with $n \leqslant (m+1)/2$. Then, for any $b_1, \dots, b_n \in \mathbb{Z}$, there exists a permutation σ on $\{1, \dots, n\}$ such that $1 + b_{\sigma(1)}, \dots, n + b_{\sigma(n)}$ are pairwise distinct modulo m.*

Proof. Observe that $m/2 > n-1 = \max_{1 \leqslant i \leqslant j \leqslant n}(j-i)$. Applying Theorem 1.3 with $\alpha_1 = \cdots = \alpha_n = 1$ and $A_1 = \cdots = A_n = \{1, \dots, n\}$, we find that there exists a permutation τ on $\{1, \dots, n\}$ such that $\tau(1)+b_1, \dots, \tau(n)+b_n$ are pairwise distinct modulo m. So the desired result follows. □

Remark 1.3. In [KS] Corollary 1.1 was applied to tree embeddings. Snevily [S] even conjectured that the condition $n \leqslant (m+1)/2$ in Corollary 1.1 can be weakened by $n < m$.

Let $G = \{a_1, \dots, a_n\}$ be an additive abelian group of order n, and let $b_1, \dots, b_n$ be elements of G with $b_1 + \cdots + b_n = 0$. In 1952 M. Hall [H] proved that there exists a permutation σ on $\{1, \dots, n\}$ such that $a_1 + b_{\sigma(1)}, \dots, a_n + b_{\sigma(n)}$ are pairwise distinct. Let σ be a permutation on $\{1, \dots, n\}$ such that $b_1 - a_{\sigma(1)}, \dots, b_n - a_{\sigma(n)}$ are pairwise distinct. Assume that $n > 1$ and $a_n = b_n = 0$. Then there exists a permutation σ' on $\{1, \dots, n-1\}$ such that

$$a_{\sigma'(i)} = a_{\sigma(i)} - a_{\sigma(n)} \neq 0 \quad \text{for every } i = 1, \dots, n-1.$$

Since $\{b_i - a_{\sigma'(i)} : i = 1, \dots, n-1\} = G \setminus \{0\}$, there is a permutation τ on $\{1, \dots, n-1\}$ such that for any $i = 1, \dots, n-1$ we have $b_i - a_{\sigma'(i)} = a_{\tau(i)}$ and hence $b_i = a_{\sigma'(i)} + a_{\tau(i)}$. In the case $G = \mathbb{Z}/n\mathbb{Z}$, this provides a positive answer to an open question of Parker (cf. [G]).

2. RELATIONS AMONG COEFFICIENTS OF CERTAIN POLYNOMIALS

As usual we let $(x)_0 = 1$ and $(x)_n = x(x-1)\cdots(x-n+1)$ for $n = 1, 2, 3, \dots$. For a polynomial

$$P(x_1, \dots, x_n) = \sum_{j_1, \dots, j_n} a_{j_1, \dots, j_n} x_1^{j_1} \cdots x_n^{j_n}$$

over a commutative ring, we write $[x_1^{j_1}\cdots x_n^{j_n}]P(x_1,\dots,x_n)$ to denote the coefficient $a_{j_1,\dots,j_n}$.

Lemma 2.1. *Let*

$$P(x_1,\dots,x_n)=\sum_{\substack{j_1,\dots,j_n\geqslant 0\\ j_1+\cdots+j_n=m}} c_{j_1,\dots,j_n}x_1^{j_1}\cdots x_n^{j_n}\in\mathbb{C}[x_1,\dots,x_n]$$

and

$$P^*(x_1,\dots,x_n)=\sum_{\substack{j_1,\dots,j_n\geqslant 0\\ j_1+\cdots+j_n=m}} c_{j_1,\dots,j_n}(x_1)_{j_1}\cdots(x_n)_{j_n}.$$

Suppose that $0\leqslant\deg P\leqslant k_1+\cdots+k_n$ *where* $k_1,\dots,k_n$ *are nonnegative integers. Then*

$$[x_1^{k_1}\cdots x_n^{k_n}]P(x_1,\dots,x_n)(x_1+\cdots+x_n)^{k_1+\cdots+k_n-\deg P}$$

coincides with

$$\frac{(\sum_{i=1}^n k_i-\deg P)!}{k_1!\cdots k_n!}P^*(k_1,\dots,k_n).$$

Proof. Let $K=k_1+\cdots+k_n-\deg P$. Then

$$\begin{aligned}&[x_1^{k_1}\cdots x_n^{k_n}]P(x_1,\dots,x_n)(x_1+\cdots+x_n)^K\\ =&[x_1^{k_1}\cdots x_n^{k_n}]\sum_{\substack{j_1,\dots,j_n\geqslant 0\\ j_1+\cdots+j_n=m}} c_{j_1,\dots,j_n}x_1^{j_1}\cdots x_n^{j_n}\sum_{\substack{i_1,\dots,i_n\geqslant 0\\ i_1+\cdots+i_n=K}} K!\frac{x_1^{i_1}\cdots x_n^{i_n}}{i_1!\cdots i_n!}\\ =&K!\sum_{\substack{j_1,\dots,j_n\geqslant 0\\ j_1+\cdots+j_n=m}} c_{j_1,\dots,j_n}\frac{(k_1)_{j_1}\cdots(k_n)_{j_n}}{k_1!\cdots k_n!}\\ =&\frac{K!}{k_1!\cdots k_n!}P^*(k_1,\dots,k_n).\end{aligned}$$

This concludes the proof. □

Let S_n denote the symmetric group of all permutations on $\{1,\dots,n\}$. For $\sigma\in S_n$ we let $\varepsilon(\sigma)$ be 1 or -1 according to whether σ is even or odd. For a matrix $A=(a_{ij})_{1\leqslant i,j\leqslant n}$ over a field the determinant and the permanent of A are defined by

$$\|A\|=\sum_{\sigma\in S_n}\varepsilon(\sigma)\prod_{i=1}^n a_{i,\sigma(i)}\quad\text{and}\quad \mathrm{per}(A)=\sum_{\sigma\in S_n}\prod_{i=1}^n a_{i,\sigma(i)}$$

respectively.

Lemma 2.1 is very useful. For example, in view of Lemmas 1.1 and 2.1, result (iii) follows from the following simple observation:

$$\begin{aligned}\|x_j^{i-1}\|^*_{1\leqslant i,j\leqslant n}=&\sum_{\sigma\in S_n}\varepsilon(\sigma)\prod_{i=1}^n(x_{\sigma(i)})_{i-1}\\ =&\|(x_j)_{i-1}\|_{1\leqslant i,j\leqslant n}=\|x_j^{i-1}\|_{1\leqslant i,j\leqslant n},\end{aligned}\tag{2.1}$$

where in the last step we note that x^r $(0\leqslant r<n)$ can be written as a linear combination of $(x)_0,\ldots,(x)_r$.

Now we present our main technique concerning the operator $P\mapsto P^*$.

Theorem 2.1. *Let $m_1,\ldots,m_n$ be nonnegative integers, and let $A=(a_{ij})_{1\leqslant i,j\leqslant n}$ be a matrix over $\mathbb{C}$. Set*

$$f(x_1,\ldots,x_n)=\|a_{ij}x_j^{m_i}\|_{1\leqslant i,j\leqslant n}P(x_1,\ldots,x_n)$$

where $P(x_1,\ldots,x_n)\in\mathbb{C}[x_1,\ldots,x_n]$ is homogeneous and

$$P(x_1,\ldots,x_{i-1},x_j,x_{i+1},\ldots,x_{j-1},x_i,x_{j+1},\ldots,x_n)=\nu P(x_1,\ldots,x_n)$$

for all $1\leqslant i<j\leqslant n$ with $\nu\in\{1,-1\}$. Then

$$f^*(x,\ldots,x)=P^*(x-m_1,\ldots,x-m_n)\prod_{i=1}^n(x)_{m_i}\times\begin{cases}\|A\| & \text{if } \nu=1,\\ \mathrm{per}(A) & \text{if } \nu=-1.\end{cases}$$

Proof. Any $\sigma\in S_n$ can be written as a product of transpositions:

$$\sigma=(i_1j_1)\cdots(i_rj_r)\ \text{ where } 1\leqslant i_s<j_s\leqslant n \text{ for } s=1,\ldots,r.$$

Thus

$$P(x_{\sigma(1)},\ldots,x_{\sigma(n)})=\nu^rP(x_1,\ldots,x_n)=\varepsilon(\sigma)^{(1-\nu)/2}P(x_1,\ldots,x_n).$$

Write $P(x_1,\ldots,x_n)=\sum_{j_1,\ldots,j_n}c_{j_1,\ldots,j_n}x_1^{j_1}\cdots x_n^{j_n}$. Then

$$\begin{aligned}&f(x_1,\ldots,x_n)\\ =&\sum_{\sigma\in S_n}\varepsilon(\sigma)\prod_{i=1}^n\left(a_{i,\sigma(i)}x_{\sigma(i)}^{m_i}\right)\times P(x_1,\ldots,x_n)\\ =&\sum_{\sigma\in S_n}\left(\varepsilon(\sigma)\prod_{i=1}^n\left(a_{i,\sigma(i)}x_{\sigma(i)}^{m_i}\right)\times\varepsilon(\sigma)^{(\nu-1)/2}P(x_{\sigma(1)},\ldots,x_{\sigma(n)})\right)\\ =&\sum_{\sigma\in S_n}\left(\varepsilon(\sigma)^{(\nu+1)/2}\prod_{i=1}^na_{i,\sigma(i)}\times\sum_{j_1,\ldots,j_n}c_{j_1,\ldots,j_n}\prod_{i=1}^nx_{\sigma(i)}^{m_i+j_i}\right).\end{aligned}$$

Therefore

$$\begin{aligned}f^*(x,\dots,x)=&\sum_{\sigma\in S_n}\varepsilon(\sigma)^{(\nu+1)/2}\prod_{i=1}^n a_{i,\sigma(i)}\times\sum_{j_1,\dots,j_n}c_{j_1,\dots,j_n}\prod_{i=1}^n(x)_{m_i+j_i}\\ =&a\sum_{j_1,\dots,j_n}c_{j_1,\dots,j_n}\prod_{i=1}^n(x)_{m_i}\times\prod_{i=1}^n(x-m_i)_{j_i}\\ =&a\prod_{i=1}^n(x)_{m_i}\times P^*(x-m_1,\dots,x-m_n)\end{aligned}$$

where

$$a=\sum_{\sigma\in S_n}\varepsilon(\sigma)^{(\nu+1)/2}\prod_{i=1}^n a_{i,\sigma(i)}=\begin{cases}\|A\| & \text{if } \nu=1,\\ \mathrm{per}(A) & \text{if } \nu=-1.\end{cases}$$

This concludes the proof. □

Corollary 2.1. *Let $m_1,\dots,m_n$ be nonnegative integers.*

(i) (Sun [Su]) *If $A=(a_{ij})_{1\leqslant i,j\leqslant n}$ is a matrix with $a_{ij}\in\mathbb{C}$, then*

$$\begin{aligned}&\Big(\|a_{ij}x_j^{m_i}\|_{1\leqslant i,j\leqslant n}\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^\delta\Big)^*(x,\dots,x)\\ &=\prod_{1\leqslant i<j\leqslant n}(m_i-m_j)^\delta\times\prod_{i=1}^n(x)_{m_i}\times\begin{cases}\|A\| & \textit{if } \delta=0,\\ \mathrm{per}(A) & \textit{if } \delta=1.\end{cases}\end{aligned}\tag{2.2}$$

(ii) We have

$$\|x_j^{m_i}\|^*_{1\leqslant i,j\leqslant n}(x-n+1,\dots,x)=\prod_{1\leqslant i<j\leqslant n}(m_j-m_i)\times\prod_{i=1}^n\frac{(x)_{m_i}}{(x)_{i-1}}.\tag{2.3}$$

Proof. As $\|x_j^{i-1}\|_{1\leqslant i,j\leqslant n}=\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)$ (Vandermonde), by (2.1) we have

$$\begin{aligned}&\Big(\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^\delta\Big)^*(x-m_1,\dots,x-m_n)\\ &=\prod_{1\leqslant i<j\leqslant n}(x-m_j-(x-m_i))^\delta=\prod_{1\leqslant i<j\leqslant n}(m_i-m_j)^\delta.\end{aligned}$$

In view of this, Theorem 2.1 yields (2.2) immediately.

By Theorem 2.1,

$$\begin{aligned}&\left(\|x_j^{n-i}\|_{1\leqslant i,j\leqslant n}\times\|x_j^{m_i}\|_{1\leqslant i,j\leqslant n}\right)^*(x,\ldots,x)\\=&n!\prod_{i=1}^n(x)_{n-i}\times\|x_j^{m_i}\|^*_{1\leqslant i,j\leqslant n}(x-n+1,\ldots,x).\end{aligned}$$

On the other hand, by part (i) we have

$$\begin{aligned}&\left(\|x_j^{n-i}\|_{1\leqslant i,j\leqslant n}\times\|x_j^{m_i}\|_{1\leqslant i,j\leqslant n}\right)^*(x,\ldots,x)\\=&(-1)^{\binom n2}\Bigg(\|x_j^{m_i}\|_{1\leqslant i,j\leqslant n}\times\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)\Bigg)^*(x,\ldots,x)\\=&n!\prod_{1\leqslant i<j\leqslant n}(m_j-m_i)\times\prod_{i=1}^n(x)_{m_i}.\end{aligned}$$

So (2.3) follows.

The proof of Corollary 2.1 is now complete. □

Remark 2.1. When $m_i=(i-1)m$ for $i=1,\ldots,n$, Corollary 2.1(ii) yields the following result related to [LS]:

$$\begin{aligned}&\Bigg(\prod_{1\leqslant i<j\leqslant n}(x_j^m-x_i^m)\Bigg)^*(x-n+1,\ldots,x)\\=&1!2!\cdots(n-1)!m^{n(n-1)/2}\frac{(x)_0(x)_m\cdots(x)_{(n-1)m}}{(x)_0(x)_1\cdots(x)_{n-1}}.\end{aligned}\tag{2.4}$$

Theorem 2.2. *Let m be any positive integer, and let $a_1,\ldots,a_n$ be complex numbers. Then*

$$\begin{aligned}&\Bigg(\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^{2m-1}\Bigg)^*(x-n+1,\ldots,x)\\=&(-1)^{(m-1)\binom n2}\frac{m!(2m)!\cdots(nm)!}{(m!)^nn!}\times\frac{(x)_0(x)_m\cdots(x)_{(n-1)m}}{(x)_0(x)_1\cdots(x)_{n-1}}\end{aligned}\tag{2.5}$$

and

$$\begin{aligned}&\Bigg(\prod_{1\leqslant i<j\leqslant n}(a_jx_j-a_ix_i)(x_j-x_i)^{2m-1}\Bigg)^*(x,\ldots,x)\\=&(-1)^{m\binom n2}\frac{m!(2m)!\cdots(nm)!}{(m!)^nn!}\mathrm{per}(a_j^{i-1})_{1\leqslant i,j\leqslant n}\prod_{r=0}^{n-1}(x)_{rm}.\end{aligned}\tag{2.6}$$

Proof. Let $P_h(x_1,\ldots,x_n)=\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^h$ for $h=1,2,3,\ldots$. In light of Theorem 2.1,

$$\begin{aligned}P_{2m}^*(x,\ldots,x)=&\left((-1)^{\binom n2}\|x_j^{n-i}\|_{1\leqslant i,j\leqslant n}P_{2m-1}(x_1,\ldots,x_n)\right)^*(x,\ldots,x)\\=&(-1)^{\binom n2}n!\prod_{i=1}^n(x)_{n-i}\times P_{2m-1}^*(x-n+1,\ldots,x)\end{aligned}$$

and

$$\begin{aligned}&\left(\|a_j^{i-1}x_j^{i-1}\|_{1\leqslant i,j\leqslant n}P_{2m-1}(x_1,\ldots,x_n)\right)^*(x,\ldots,x)\\=&\frac{\operatorname{per}(a_j^{i-1})_{1\leqslant i,j\leqslant n}}{\operatorname{per}(a_j^0)_{1\leqslant i,j\leqslant n}}\left(\|a_j^0x_j^{i-1}\|_{1\leqslant i,j\leqslant n}P_{2m-1}(x_1,\ldots,x_n)\right)^*(x,\ldots,x)\\=&\frac{\operatorname{per}(a_j^{i-1})_{1\leqslant i,j\leqslant n}}{n!}P_{2m}^*(x,\ldots,x).\end{aligned}$$

By Theorem 3.1 of Hou and Sun [HS],

$$P_{2m}^*(x,\ldots,x)=(-1)^{m\binom n2}\frac{m!(2m)!\cdots(nm)!}{(m!)^n}(x)_0(x)_m\cdots(x)_{(n-1)m}.$$

So we have the desired (2.5) and (2.6). □

Corollary 2.2. *Let k,m,n be positive integers with $k>m(n-1)$. Then*

$$\begin{aligned}&[x_1^{k-n}\cdots x_n^{k-1}](x_1+\cdots+x_n)^{(k-1)n-mn(n-1)}\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^{2m-1}\\&=(-1)^{(m-1)\binom n2}\frac{m!(2m)!\cdots(nm)!}{(m!)^nn!}\cdot\frac{((k-1-m(n-1))n)!}{\prod_{r=0}^{n-1}(k-1-rm)!}.\end{aligned}\tag{2.7}$$

In particular,

$$\left[\prod_{i=1}^n x_i^{(m-1)(n-1)+i-1}\right]\prod_{1\leqslant i<j\leqslant n}(x_j-x_i)^{2m-1}=(-1)^{(m-1)\binom n2}\frac{(mn)!}{(m!)^nn!}.\tag{2.8}$$

Proof. Combining Lemma 2.1 with (2.5) we obtain (2.7). (2.8) follows from (2.7) in the case $k=m(n-1)+1$. □

Remark 2.3. Let $m_1,\ldots,m_n$ be nonnegative integers. A confirmed conjecture of Dyson [D] can be stated as follows:

$$\begin{aligned}&[x_1^{m_1(n-1)}\cdots x_n^{m_n(n-1)}]\prod_{1\leqslant i<j\leqslant n}(x_i-x_j)^{m_i+m_j}\\&=(-1)^{\sum_{j=1}^n(j-1)m_j}\frac{(m_1+\cdots+m_n)!}{m_1!\cdots m_n!}.\end{aligned}$$

(See, e.g., Zeilberger [Z].) Compared with this deep result, our (2.8) seems interesting too.

3. Proofs of Theorems 1.1–1.3

Proof of Theorem 1.1. The case $n = 1$ or $k-1 < m(n-1)$ is trivial. Below we assume $n \geqslant 2$ and $l = k-1-m(n-1) \geqslant 0$.

As $|F| \geqslant p > mn \geqslant 2m$ we can extend each S_{ij} $(1 \leqslant i < j \leqslant n)$ to a subset S_{ij}^* of F with cardinality $2m-1$. By Lemma 1.1 it suffices to show that

$$[x_1^{k-n} \cdots x_n^{k-1}](x_1 + \cdots + x_n)^{ln} \prod_{1 \leqslant i < j \leqslant n} \prod_{c \in S_{ij}^*} (x_j - x_i + c)$$

does not vanish. Let e denote the multiplicative identity of the field F. Then the above coefficient equals he where

$$h = [x_1^{k-n} \cdots x_n^{k-1}](x_1 + \cdots + x_n)^{ln} \prod_{1 \leqslant i < j \leqslant n} (x_j - x_i)^{2m-1} \in \mathbb{Z}.$$

By Corollary 2.2,

$$h = (-1)^{(m-1)\binom{n}{2}} \frac{m!(2m)! \cdots (nm)!}{(m!)^n n!} \cdot \frac{(ln)!}{\prod_{r=0}^{n-1}(k-1-rm)!}.$$

As $p > mn$ and $p > ln$, p does not divide h and hence $he \neq 0$. This concludes the proof. $\square$

Proof of Theorem 1.2. To avoid triviality, we assume $n \geqslant 2$ and $l = k-1-m(n-1) \geqslant 0$. As q is odd, the norms of those $1 - \zeta_s/\zeta_t$ $(1 \leqslant s < t \leqslant n)$ (with respect to the field extension $\mathbb{Q}(e^{2\pi i/q})/\mathbb{Q}$) are odd integers and hence $\|\zeta_t^{s-1}\|_{1 \leqslant s,t \leqslant n} = \prod_{1 \leqslant s < t \leqslant n}(\zeta_t - \zeta_s)$ is not an algebraic integer times two. Therefore $\operatorname{per}(\zeta_t^{s-1})_{1 \leqslant s,t \leqslant n} \neq 0$ as observed by Dasgupta et al. [DKSS]. By Lemma 2.1 and (2.6),

$$[x_1^{k-1} \cdots x_n^{k-1}](x_1 + \cdots + x_n)^{ln} \prod_{1 \leqslant s < t \leqslant n} (\zeta_t x_t - \zeta_s x_s)(x_t - x_s)^{2m-1} \neq 0.$$

Applying Lemma 1.1 we then obtain the desired result. $\square$

Proof of Theorem 1.3. For $1 \leqslant i < j \leqslant n$, let r_{ij} denote the unique integer in the interval $(-m_{ij}/2, m_{ij}/2]$ which is congruent to $b_i - b_j$ modulo m_{ij}. For $x_i \in A_i$ and $x_j \in A_j$, as $|x_i - x_j| < m_{ij}/2$ we have

$$x_i + b_i \equiv x_j + b_j \pmod{m_{ij}} \iff x_j - x_i = r_{ij}.$$

Note also that

$$\operatorname{per}(\alpha_j^{i-1})_{1 \leqslant i,j \leqslant n} = \sum_{\sigma \in S_n} \prod_{i=1}^n \alpha_{\sigma(i)}^{i-1} > 0.$$

Thus Theorem 1.3 follows from (2.6) and Lemmas 1.1 and 2.1. □

Acknowledgments. The main part of this work was done during the first author's visit to the second author's institute, Z. W. Sun would like to thank the Institute of Mathematics, Academia Sinica (Taiwan) for its support. The paper was revised during Sun's visit to the University of California at Irvine, he is indebted to Prof. Daqing Wan for the invitation.

References

[A1] N. Alon, *Combinatorial Nullstellensatz*, Combin. Probab. Comput. **8** (1999), 7–29.

[A2] N. Alon, *Additive Latin transversals*, Israel J. Math. **117** (2000), 125–130.

[A3] N. Alon, *Discrete mathematics: methods and challenges*, in: Proceedings of the International Congress of Mathematicians (Beijing, 2002), Vol. I, Higher Ed. Press, Beijing, 2002, pp. 119–135.

[ANR1] N. Alon, M. B. Nathanson and I. Z. Ruzsa, *Adding distinct congruence classes modulo a prime*, Amer. Math. Monthly **102** (1995), 250–255.

[ANR2] N. Alon, M. B. Nathanson and I. Z. Ruzsa, *The polynomial method and restricted sums of congruence classes*, J. Number Theory **56** (1996), 404–417.

[AT] N. Alon and M. Tarsi, *A nowhere-zero point in linear mappings*, Combinatorica **9** (1989), 393–395.

[DKSS] S. Dasgupta, G. Károlyi, O. Serra and B. Szegedy, *Transversals of additive Latin squares*, Israel J. Math. **126** (2001), 17–28.

[DH] J. A. Dias da Silva and Y. O. Hamidoune, *Cyclic spaces for Grassmann derivatives and additive theory*, Bull. London Math. Soc. **26** (1994), 140–146.

[D] F. J. Dyson, *Statistical theory of the energy levels of complex systems I*, J. Math. Phys. **3** (1962), 140–156.

[G] R. K. Guy, *Parker's permutation problem involves the Catalan numbers*, Amer. Math. Monthly **100** (1993), 287–289.

[H] M. Hall, *A combinatorial problem on abelian groups*, Proc. Amer. Math. Soc. **3** (1952), 584–587.

[HS] Q. H. Hou and Z. W. Sun, *Restricted sums in a field*, Acta Arith. **102** (2002), 239–249.

[KS] A. E. Kézdy and H. S. Snevily, *Distinct sums modulo n and tree embeddings*, Combin. Probab. Comput. **11** (2002), 35–42.

[LS] J. X. Liu and Z. W. Sun, *Sums of subsets with polynomial restrictions*, J. Number Theory **97** (2002), 301–304.

[N] M. B. Nathanson, *Additive Number Theory: Inverse Problems and the Geometry of Sumsets* (Graduate texts in mathematics; 165), Springer, New York, 1996.

[S] H. S. Snevily, *The Cayley addition table of $\mathbb{Z}_n$*, Amer. Math. Monthly **106** (1999), 584–585.

[Su] Z. W. Sun, *On Snevily's conjecture and restricted sumsets*, J. Combin. Theory Ser. A **103** (2003), 291–304.

[Z] D. Zeilberger, *A combinatorial proof of Dyson's conjecture*, Discrete Math. **41** (1982), 317–321.